

# **Challenges of convection-permitting regional climate simulations for future climate projection in Japan**

**- Program for Risk Information on Climate Change, SOUSEI program -**

NCAR  
Sep. 7, 2016

**Hiroaki Kawase**

(Meteorological Research Institute [MRI], Tsukuba, Japan)

Hidetaka Sasaki, Akihiko Murata, Masaya Nosaka, Izuru Takayabu (MRI)



Toshinori Aoyagi (JMA), Rui Ito (NIED)

# CONTENTS

- 1. Introduction of Japanese large project of climate projection: SOSEI program**
- 2. Methodology of global and regional climate simulations in SOUSE program**
- 3. Impact of high-resolution simulation on precipitation and snow depth in Japan**
- 4. Summary**

# Introduction

- Since there are many mountain ranges in Japan, the local climate is so complicated.
- The urban areas are locally influenced by the heat island effects.
- The high resolution climate simulations are required to simulate the mesoscale heavy rainfall/snowfall, tropical cyclones, urban climate, and orographic effects in Japan.





# Introduction

Japanese large scientific program, **Program for Risk Information on Climate Change**, called as **SOUSEI program**, conducts present and future climate simulations to generate basic information required for managing various risks resulting from climate change.



## Four themes in SOUSEI program

<http://www.jamstec.go.jp/sousei/eng/index.html>



Using the world-class supercomputers, such as Earth Simulator, we are pursuing research and development in which all themes are organically linked. Our research and development include prediction and diagnosis of imminent global climate change expected to occur within a few years or decades, research on greenhouse gas emission scenarios and associated long-term climate change projections, development of probabilistic climate change projection techniques, and development of technology for precise impact assessment, etc.

Our team

### Theme C

**Development of basic technology for risk information on climate change (= Downscaling)**



**Theme A** Prediction and diagnosis of imminent global climate change

- Understanding mechanisms of climate variability and change
- Development of an integrated prediction system for global climate studies

**Theme B** Climate change projection contributing to stabilization target setting

- Long-term global change projection based on diverse scenarios
- Obtaining scientific perceptions on large-scale variations and modifications of climate

**Theme C** Development of basic technology for risk information on climate change

- Probabilistic climate prediction for risk assessment
- Producing a standard climate scenario by using super high resolution models

**Theme D** Precise impact assessments on climate change




- Climate change impacts on natural hazards
- Climate change impacts on water resources
- Climate change impacts on ecosystem and biodiversity

# Theme C: Development of basic technology for risk information on climate change


University of Tsukuba, NIED, ISM, AORI, ISEE, MRI

Representative: I. Takayabu @ MRI

## (i) Estimate PDF of extreme events



## (ii) Find the worst case scenario





# Theme C: Development of basic technology for risk information on climate change

University of Tsukuba, NIED, ISM, AORI, ISEE, MRI

Representative: I. Takayabu @ MRI

## (i) Estimate PDF of extreme events



## (ii) Find the worst case scenario


- In this presentation, I will introduce the approach of present and future climate simulation using high-resolution global and regional climate models.
- Then, I will show the biases of precipitation in Japan and impacts of high-resolution experiments on precipitation and snow depth in Japan.  
(Unfortunately, the future projections with 2km grid spacing is now calculating.)




# Methodology of Climate Experiment in SOUSEI theme C

20km MRI-AGCM\*1  
and 2km/5km NHRCM\*2


\*MRI-AGCM: Meteorological Research Institute AGCM  
\* NHRCM: Non-Hydrostatic Regional Climate Model



2km NHRCM



5km NHRCM



20km MRI-AGCM

High resolution AMIP-type present climate simulation and future climate projection. [Mizuta et al. 2012]




## Past simulation and future projection by MRI-AGCM [Mizuta et al. 2012]


AMIP type time-slice simulations

Present climate: 1980-2000 (SST: HadISST)

Future climate: 2076-2096 (SST: RCP8.5)\*




\* Future climate simulations were performed using three SSTs anomaly in CMIP5 models under RCP8.5. To obtain three patterns of SST change, a cluster analysis was conducted.




**NCAR type**

(Nor



**HadGEM2 type**  
(including MRI-CGCM3)



**GFDL type**

**ensemble mean**



# Methodology of Climate Experiment in SOUSEI

|                       |   |  |
|-----------------------|---|--|
| <b>Model</b>          | <b>Non-Hydrostatic Regional Climate Model (NHRCM)</b><br>(A fully compressible and nonhydrostatic model) [Sasaki et al. 2011] |  |
| <b>Grid spacing</b>   | <b>5km</b>  | <b>2km</b>   |
| <b>Grid number</b>    | 527x804x50  | 525x1721x60  |
| <b>Microphysics</b>   | Bulk-type cloud microphysics [Ikawa et al., 1994]   |  |
| <b>Convection</b>     | <i>Kain and Fritsch (1993)</i>  | -  |
| <b>Radiation</b>      | Clear-sky [Yabu et al., 2005], Cloudy [Kitagawa et al., 2000]   |  |
| <b>Boundary layer</b> | Improved MYNN Level 3 [Nakanishi and Niino, 2004]   |  |
| <b>Land surface</b>   | Improved MRI/JMA SiB [Hirai and Ohizumi, 2004]  |  |
| <b>Integration</b>    | 20 years [Jul. 20 – Aug. 31 (following year) in each year]  |  |
| <b>Urban</b>          | -   | Square Prism Urban Canopy (SPUC)<br>[Aoyagi and Seino, 2011] |


# SPUC (Urban canopy model)

## Difference of temperature between 2006 land use and 1976 land use [lateral boundary condition: 2006]


summer

winter



EXP2006HR - EXP1976, JUL\_AUG



EXP2006HR - EXP1976, JAN\_FEB



differential [°C]





Higher temperature is simulated in both season  
around Tokyo Metropolitan area.



[Aoyagi and Seino, 2012]

# Precipitation biases



## Biases of annual precipitation in each region



Biases and RMSE of annual precipitation in 2km NHRCM are less than those in 5km NHRCM.









## Relative frequency distribution

Murata et al. (2015)

**DAILY** precipitation




**HOURLY** precipitation



The relative frequency distribution of daily precipitation is well reproduced by 2km NHRCM than by 5km NHRCM when compared with the observed precipitation data except for higher intensities.



# Simulated snowfall and snow depth

## 20 years mean maximum snow depth [NHRCM] – [Observation]




# Simulated snowfall in the mountainous areas


## Seasonal variation of snow depth



### ■ Bijyodaira site (977m)



### ■ Midagahara site (1,930m)






# Simulated snowfall and snow depth

## East Asian Winter Monsoon




From JMA web site




Colder air from Siberia and warmer SST result in more developed snow clouds and heavier snowfall

# Simulated snowfall and snow depth

## East Asian Winter Monsoon




From JMA web site




- The major differences between 2km and 5km NHRCMs are
  - » Highly resolved topography
  - » With/without cumulus parameterization
- ➔ Sensitivity experiments on 2014/15 winter with 1km and 5km. Checking horizontal distribution of precipitation

# Simulated snowfall in the mountainous areas

## Winter total precipitation



## Comparison of precipitation with 1kmNHRCM





# Summary

## ■ SOUSEI program

- The **SOUSEI program (theme C)** performs present and future climate simulations using **MRI-AGCM** and **NHRCM**.
- The NHRCM with 2km grid-spacing are performed without cumulus parameterization and with urban canopy model (SPUC).
- **The 2km NHRCM better simulated the precipitation and maximum snow depth** in Japan rather than 5km NHRCM did.

## ■ Resolution dependency of snowfall and rainfall

- **Annual maximum snow depth is overestimated by the NHRCM with 5km** in the mountainous areas, which results from overestimation of winter precipitation.
- **High-resolution experiments without KF scheme** is important to accurately simulate Japanese snowfall.

*Thank you for your attention*

